arXiv:1209.4632v3 [math.FA] 22 Nov 2012

THE BOHNENBLUST-HILLE INEQUALITY FOR REAL HOMOGENEOUS POLYNOMIALS IS HYPERCONTRACTIVE AND THIS RESULT IS OPTIMAL

J. R. CAMPOS, G.A. MUÑOZ-FERNÁNDEZ, D. PELLEGRINO, AND J.B. SEOANE-SEPÚLVEDA

Abstract. It was recently shown that the optimal constants satisfying the multilinear Bohnenblust–Hille inequalities (real and complex) have a subpolynomial growth; as a consequence, A. Montanaro provided striking applications of this fact, for the case of real scalars, to Quantum Information Theory. In this paper, we show that contrary to the case of multilinear mappings, the optimal constants for the polynomial Bohnenblust–Hille inequality for real scalars do not have a (sub)polynomial growth. In fact, as the title of this paper states, we show that the polynomial Bohnenblust–Hille inequality (for real scalars) is hypercontractive and, moreover, this result cannot be improved. For complex scalars the hypercontractivity of the polynomial Bohnenblust–Hille inequality was recently proved by Defant, Frerick, Ortega-Cerdá, Ounaïes, and Seip (*Annals of Mathematics*, 2011). The optimality of this result for complex scalars, however, remains still open. Our proof of the hypercontractivity is presented in a simple form, employing an old result on complexifications of polynomials. We also introduce a technique, which is a combination of the Krein–Milman Theorem together with a description of the geometry of the unit ball of polynomial spaces on ℓ_∞^2, to generate lower bounds for the constants in the Bohnenblust–Hille inequality.

1. The polynomial Bohnenblust–Hille inequality

1.1. **Complex Scalars.** The polynomial Bohnenblust–Hille inequality appeared in 1931 [5] as a crucial tool for solving the famous Bohr's radius problem (see, also, [3, 4, 11–13]). This inequality asserts that there is a constant $D_{\mathbb{C},m} \geq 1$ so that

$$\left(\sum_{|\alpha|=m} |a_\alpha|^{\frac{2m}{m+1}} \right)^{\frac{m+1}{2m}} \leq D_{\mathbb{C},m} \, \|P\| \tag{1.1}$$

for all m-homogeneous polynomials $P : \ell_\infty^N (\mathbb{C}) \to \mathbb{C}$ given by

$$P(z) = \sum_{|\alpha|=m} a_\alpha z^\alpha .$$

The original estimates of $D_{\mathbb{C},m}$ presented a very "bad growth" (of order m^m) and only very recently Defant, Frerick, Ortega-Cerdá, Ounaïes and Seip [9] proved that this inequality is hypercontractive:

$$D_{\mathbb{C},m} \leq \left(1 + \frac{1}{m-1}\right)^{m-1} \sqrt{m} \left(\sqrt{2}\right)^{m-1} \quad \text{for } m \geq 2. \tag{1.2}$$

2010 *Mathematics Subject Classification.* 46G25, 47L22, 47H60.

Key words and phrases. Bohnenblust–Hille inequality, Absolutely summing operators, Krein–Milman Theorem.

D. Pellegrino was supported by CNPq Grants 301237/2009-3, 477124/2012-7, INCT-Matemática and CAPES-NF. G.A. Muñoz-Fernández and J. B. Seoane-Sepúlveda were supported by MTM2009-07848.

The above result has several specific applications in Harmonic Analysis and Dirichlet Series, as shown in [9]. The visible lack of interest in the real scalar version of the Bohnenblust–Hille inequality was probably due to the lack of potential applications for the last 80 years.

1.2. When Quantum Information Theory comes into play.

Let $\mathbb{K}$ be the real or complex scalar field. The multilinear Bohnenblust–Hille inequality asserts that there is a constant $C_{\mathbb{K},n}$ (depending only on n and $\mathbb{K}$) such that

$$\left(\sum_{i_1,\ldots,i_m=1}^{N}\left|A(e_{i_1},\ldots,e_{i_n})\right|^{\frac{2n}{n+1}}\right)^{\frac{n+1}{2n}}\leq C_{\mathbb{K},n}\left\|A\right\|$$

for all continuous n-linear forms A on $\ell_\infty^N(\mathbb{K})\times\cdots\times\ell_\infty^N(\mathbb{K})$.

The Bohnenblust–Hille inequalities (multilinear and polynomial) for real scalars have recently been investigated thoroughly (see, e.g., [14, 17, 18, 26, 30–32]). For example, it was proved that for the multilinear case the optimal constants of the Bohnenblust–Hille inequality have a subpolynomial growth (for both real and complex scalars) [31].

In 2012, the panorama of applications changed when A. Montanaro [25] provided explicit applications to Quantum Information Theory of the hypercontractivity of the constants of the multilinear Bohnenblust–Hille inequality for real scalars. Montanaro's work shows, contrary to what was believed in the past, that the study of the growth of the constants in the case of real scalars also has crucial applications.

1.3. Summary of the results.

In general, the constants for the case of real scalars seem to be bigger (although this result is still not known) than the constants for the complex case (see [32]). Thus, the hypercontractivity of the polynomial Bohnenblust–Hille inequality is not, at all, any direct consequence of the case of complex scalars.

One of the goals of this paper is to stress the fact that the polynomial Bohnenblust–Hille inequality for real scalars is also hypercontractive. Moreover, we also show that this hypercontractive growth cannot be improved in any possible way. Also, we shall introduce a technique (which is a combination of the Krein–Milman Theorem together with a description of the geometry of the unit ball of polynomial spaces on ℓ_∞^2) to generate lower bounds for the constants in the Bohnenblust–Hille inequality. A combination of the above results provides a very good estimate for the hypercontractivity constant of the polynomial Bohnenblust–Hille inequality for real scalars. In this direction we show that the hypercontractivity constant $H_{\mathbb{R}}$ of the polynomial Bohnenblust–Hille inequality (for real scalars) is such that

$$1.495 < H_{\mathbb{R}} < 3.296.$$

Moreover, given any $\varepsilon > 0$, then

$$H_{\mathbb{R}} \leq 2\sqrt{2}+\varepsilon < 2.829,$$

if only sufficiently large values of m are considered.

2. The polynomial Bohnenblust–Hille inequality: The real case and "bad" constants

It is well-known that the multilinear Bohnenblust–Hille inequality holds for real scalars. In this section we present the canonical approach, which shows that the constants involved grow very rapidly. Let us introduce some notation and a few basic results.

Given $\alpha = (\alpha_1, \ldots, \alpha_n) \in (\mathbb{N}^*)^n$, define $|\alpha| := \alpha_1 + \cdots + \alpha_n$ and

$$\binom{m}{\alpha} := \frac{m!}{\alpha_1! \cdots \alpha_n!},$$

for $|\alpha| = m \in \mathbb{N}^*$. Also, $\mathbf{x}^\alpha$ stands for the monomial $x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ for $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{K}^n$. We define the space $\mathcal{P}(^m\ell_\infty^n)$ of the m homogeneous polynomials on $\mathbb{K}^n$, with $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$, as the space over $\mathbb{K}$ generated by the monomials $\mathbf{x}^\alpha$ such that $\alpha \in (\mathbb{N}^*)^n$ and $|\alpha| = m$. A standard algebraic result states that for each polynomial $P \in \mathcal{P}(^m\ell_\infty^n)$ (real or complex), there exists a unique symmetric m-linear form L on $\ell_\infty^n \times \overset{m}{\cdots} \times \ell_\infty^n$, called the polar of P, such that $P(\mathbf{x}) = L(\mathbf{x}, \overset{m}{\ldots}, \mathbf{x})$ for all $\mathbf{x} \in \mathbb{K}^n$. If $P \in \mathcal{P}(^m\ell_\infty^n)$ and L is its polar, then we define the canonical norm of P and L, respectively by

$$\begin{aligned} \|P\| &= \sup\{|P(\mathbf{x})| : \ \|\mathbf{x}\|_\infty \leq 1\}, \\ \|L\| &= \sup\{|L(\mathbf{x_1}, \ldots, \mathbf{x_m})| : \ \|\mathbf{x_k}\|_\infty \leq 1 \text{ for } k = 1, \ldots, m\}, \end{aligned}$$

where $\|\cdot\|_\infty$ is the usual ℓ_∞ norm on $\mathbb{K}^n$.

Having all this in mind, a straightforward consequence of the multinomial formula yields the following relationship between the coefficients of a homogeneous polynomial and the polar of the polynomial (this lemma appears in [10] and it is essentially *folklore*).

Lemma 2.1. *If P is a homogeneous polynomial of degree m on $\mathbb{K}^n$ given by*

$$P(x_1, \ldots, x_n) = \sum_{|\alpha|=m} a_\alpha \mathbf{x}^\alpha,$$

and L is the polar of P, then

$$L(e_1^{\alpha_1}, \ldots, e_n^{\alpha_n}) = \frac{a_\alpha}{\binom{m}{\alpha}},$$

where $\{e_1, \ldots, e_n\}$ is the canonical basis of $\mathbb{K}^n$ and $e_k^{\alpha_k}$ stands for e_k repeated α_k times.

The following proof of the polynomial Bohnenblust–Hille inequality is a special case of [15, Lemma 5], but we present it here for the sake of completeness of this paper.

Theorem 2.2 (Bohnenblust–Hille inequality: Polynomial case and constants in the mathematical "folklore")**.** *If P is a homogeneous polynomial of degree m on $\mathbb{R}^n$ given by*

$$P(x_1, \ldots, x_n) = \sum_{|\alpha|=m} a_\alpha \mathbf{x}^\alpha,$$

then

$$\left(\sum_{|\alpha|=m} |a_\alpha|^{\frac{2m}{m+1}} \right)^{\frac{m+1}{2m}} \leq D_{\mathbb{R},m} \|P\|$$

with

$$D_{\mathbb{R},m} = C_{\mathbb{R},m} \frac{m^m}{(m!)^{\frac{m+1}{2m}}}.$$

Proof. From Lemma 2.1 we have

$$\begin{aligned}\sum_{|\alpha|=m} |a_\alpha|^{\frac{2m}{m+1}} &= \sum_{|\alpha|=m} \left(\binom{m}{\alpha} |L(e_1^{\alpha_1}, \ldots, e_n^{\alpha_n})|\right)^{\frac{2m}{m+1}} \\ &= \sum_{|\alpha|=m} \binom{m}{\alpha}^{\frac{2m}{m+1}} |L(e_1^{\alpha_1}, \ldots, e_n^{\alpha_n})|^{\frac{2m}{m+1}} .\end{aligned}$$

However, for every choice of α, the term

$$|L(e_1^{\alpha_1}, \ldots, e_n^{\alpha_n})|^{\frac{2m}{m+1}}$$

is repeated $\binom{m}{\alpha}$ times in the sum

$$\sum_{i_1,\ldots,i_m=1}^{n} |L(e_{i_1}, \ldots, e_{i_m})|^{\frac{2m}{m+1}} .$$

Thus

$$\sum_{|\alpha|=m} \binom{m}{\alpha}^{\frac{2m}{m+1}} |L(e_1^{\alpha_1}, \ldots, e_n^{\alpha_n})|^{\frac{2m}{m+1}} = \sum_{i_1,\ldots,i_m=1}^{n} \binom{m}{\alpha}^{\frac{2m}{m+1}} \frac{1}{\binom{m}{\alpha}} |L(e_{i_1}, \ldots, e_{i_m})|^{\frac{2m}{m+1}}$$

and, since

$$\binom{m}{\alpha} \leq m!$$

we have

$$\sum_{|\alpha|=m} \binom{m}{\alpha}^{\frac{2m}{m+1}} |L(e_1^{\alpha_1}, \ldots, e_n^{\alpha_n})|^{\frac{2m}{m+1}} \leq (m!)^{\frac{m-1}{m+1}} \sum_{i_1,\ldots,i_m=1}^{n} |L(e_{i_1}, \ldots, e_{i_m})|^{\frac{2m}{m+1}} .$$

We finally obtain

$$\begin{aligned}\left(\sum_{|\alpha|=m} |a_\alpha|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}} &\leq \left((m!)^{\frac{m-1}{m+1}} \sum_{i_1,\ldots,i_m=1}^{n} |L(e_{i_1}, \ldots, e_{i_m})|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}} \\ &= (m!)^{\frac{m-1}{2m}} \left(\sum_{i_1,\ldots,i_m=1}^{n} |L(e_{i_1}, \ldots, e_{i_m})|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}} \\ &\leq (m!)^{\frac{m-1}{2m}} C_{\mathbb{R},m} \|L\| .\end{aligned}$$

On the other hand, it is well-known that

$$\|L\| \leq \frac{m^m}{m!} \|P\|$$

and hence

$$\begin{aligned}\left(\sum_{|\alpha|=m}|a_\alpha|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}} &\leq C_{\mathbb{R},m}\,(m!)^{\frac{m-1}{2m}}\,\frac{m^m}{m!}\,\|P\| \\ &= C_{\mathbb{R},m}\frac{m^m}{(m!)^{\frac{m+1}{2m}}}\,\|P\|\,.\end{aligned}$$

□

Even if we use the best known values for $C_{\mathbb{R},m}$, the following table shows that the estimates for $D_{\mathbb{R},m}$ are, actually, not good at all:

m	$C_{\mathbb{R},m}$ ([32], 2012)	$D_{\mathbb{R},m}$
3	≈ 1.7818	≈ 14.5698
4	2	≈ 70.2486
5	≈ 2.2974	≈ 406.048
6	≈ 2.5198	$\approx 2,532.23$
7	≈ 2.6918	$\approx 16,984.6$
8	≈ 2.8284	$\approx 121,803$
9	≈ 3.0548	$\approx 964,757$
10	≈ 3.2490	$\approx 8.01459\cdot 10^{6}$
25	≈ 5.1366	$\approx 3.63114\cdot 10^{22}$
100	≈ 10.5101	$\approx 1.76504\cdot 10^{121}$
500	≈ 23.9012	$\approx 4.85435\cdot 10^{782}$
1,000	≈ 34.4078	$\approx 2.82183\cdot 10^{1,716}$

3. The Bohnenblust–Hille inequality for real polynomials is hypercontractive

Our argument is quite simplified by the use of complexifications of polynomials. In particular we are interested in the following result due to Visser [36], which generalizes and old result of Chebyshev:

Theorem 3.1 (Visser, [36], 1946). *Let*

$$P(y_1,\ldots,y_N)=\sum_{|\alpha|\leq k}a_\alpha y_1^{\alpha_1}\cdots y_N^{\alpha_N},$$

with $\alpha=(\alpha_1,\ldots,\alpha_N)$, $|\alpha|=\alpha_1+\cdots+\alpha_N$, *be a polynomial of total degree at most* $k\in\mathbb{N}$ *in several variables* $(y_1,\ldots,y_N)$ *and with real coefficients. Suppose* P_k *is the* k*-homogeneous polynomial defined by*

$$P_k(y_1,\ldots,y_N)=\sum_{|\alpha|=k}a_\alpha y_1^{\alpha_1}\cdots y_N^{\alpha_N}.$$

Then we have

$$\max_{|z_1|=\cdots=|z_N|=1}|P_k(z_1,\ldots,z_N)|\leq 2^{k-1}\cdot\max_{x_1,\ldots,x_N\in[-1,1]}|P(x_1,\ldots,x_N)|.$$

In particular, if P *is homogeneous, then*

$$\max_{|z_1|=\cdots=|z_N|=1}|P(z_1,\ldots,z_N)|\leq 2^{k-1}\cdot\max_{x_1,\ldots,x_N\in[-1,1]}|P(x_1,\ldots,x_N)|.$$

Moreover, the constant 2^{k-1} *cannot be replaced by any smaller one.*

Let us recall that the previous result was generalized by Muñoz, Sarantopoulos and Tonge [28, Proposition 16] for polynomials on a real Banach space using any *natural complexification procedure* (see [28] for a complete account on complexifications of polynomials). Other results concerning the norm of the complex extension of a homogeneous or a non-homogeneous polynomial can be found in [1, 19, 22, 34, 35] just to mention a few.

Let $P : \ell_\infty^N(\mathbb{R}) \to \mathbb{R}$ be an m-homogeneous polynomial

$$P(x) = \sum_{|\alpha|=m} a_\alpha x^\alpha$$

and consider the complexification $P_\mathbb{C} : \ell_\infty^N(\mathbb{C}) \to \mathbb{C}$ of P given by

$$P_\mathbb{C}(z) = \sum_{|\alpha|=m} a_\alpha z^\alpha.$$

From Theorem 3.1 above we know that

$$\|P_\mathbb{C}\| \leq 2^{m-1} \|P\|.$$

We thus have, from (1.1) and (1.2),

$$\left(\sum_{|\alpha|=m} |a_\alpha|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}} \leq \left(1 + \frac{1}{m-1}\right)^{m-1} \sqrt{m} \left(\sqrt{2}\right)^{m-1} \|P_\mathbb{C}\|.$$

Therefore applying Theorem 3.1 and the facts that $\left(1+\frac{1}{k}\right)^{\frac{1}{k}} < e$ and $\frac{e\sqrt{k}}{2\sqrt{2}} < a^k$ for all $k \in \mathbb{N}$ with $a = \sqrt[3]{\frac{e\sqrt{3}}{2\sqrt{2}}}$, then

$$\begin{aligned}\left(\sum_{|\alpha|=m} |a_\alpha|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}} &\leq \left(1 + \frac{1}{m-1}\right)^{m-1} \sqrt{m} \left(2\sqrt{2}\right)^{m-1} \|P\| \\ &< \frac{e\sqrt{m}}{2\sqrt{2}} (2\sqrt{2})^m \cdot \|P\| \\ &\leq (2\sqrt{2}a)^m \|P\|.\end{aligned}$$

Observe that $2\sqrt{2}a < 3.352$. This hypercontractive estimate on the growth of $D_{\mathbb{R},m}$ can be slightly improved by considering better estimates than (1.2) for small values of m. We will do this later on.

4. HYPERCONTRACTIVITY FOR REAL SCALARS CAN NOT BE IMPROVED

The following result asserts that the hypercontractivity in the real case cannot be improved:

Theorem 4.1. *The growth of the polynomial Bohnenblust–Hille inequality for real scalars cannot be better than hypercontractive. Moreover,*

$$D_{\mathbb{R},m} > \left(\frac{2\sqrt[4]{3}}{\sqrt{5}}\right)^m > (1.177)^m$$

for all positive integers m.

Proof. Let m be an even integer. Consider the m-homogeneous polynomial

$$P_m(x_1,\ldots,x_m)=\left(x_1^2-x_2^2+x_1x_2\right)\left(x_3^2-x_4^2+x_3x_4\right)\cdots\left(x_{m-1}^2-x_m^2+x_{m-1}x_m\right).$$

Since $\|P_2\|=5/4$, it is simple to see that

$$\|P_m\|=(5/4)^{m/2}.$$

From the Bohnenblust–Hille inequality for P_m we have

$$\left(\sum_{|\alpha|=m}|a_\alpha|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}}\leq D_{\mathbb{R},m}\,\|P_m\|,$$

that is,

$$D_{\mathbb{R},m}\geq\frac{\left(3^{\frac{m}{2}}\right)^{\frac{m+1}{2m}}}{\left(\frac{5}{4}\right)^{\frac{m}{2}}}\geq\frac{\left(\sqrt{3}\right)^{\frac{m+1}{2}}}{\left(\frac{5}{4}\right)^{\frac{m}{2}}}>\left(\frac{2\sqrt[4]{3}}{\sqrt{5}}\right)^{m}.$$

Now let us suppose that m is odd. Keeping the previous notation, consider the m homogeneous polynomial

$$Q\left(x_1,\ldots,x_{2m}\right)=\left(x_{2m}+x_{2m-1}\right)P_{m-1}\left(x_1,\ldots,x_{m-1}\right)+\left(x_{2m}-x_{2m-1}\right)P_{m-1}\left(x_m,\ldots,x_{2m-2}\right).$$

So we have

$$D_{\mathbb{R},m}\geq\frac{\left(4\cdot3^{\frac{m-1}{2}}\right)^{\frac{m+1}{2m}}}{2\cdot\left(\frac{5}{4}\right)^{\frac{m-1}{2}}}>2^{m-1+\frac{1}{m}}\left(\frac{\sqrt[4]{3}}{\sqrt{5}}\right)^{m-1}>\left(\frac{2\sqrt[4]{3}}{\sqrt{5}}\right)^{m-1}.$$

□

5. Improving the lower bounds for $D_{\mathbb{R},m}$: some technical background

The following result is crucial for our goals:

Theorem 5.1 (consequence of the Krein–Milman Theorem). *If C is a convex body in a Banach space and $f:C\to\mathbb{R}$ is a convex function that attains its maximum, then there is an extreme point $e\in C$ so that $f(e)=\max\{f(x):x\in C\}$.*

This consequence of the Krein–Milman Theorem ([23]) provides good lower estimates on the constants when it is combined with a description of the geometry of the unit ball of a polynomial space on ℓ_∞^m. The problem of finding the extreme points of the unit ball of a polynomial space has been largely studied in the past few years. In particular, the following results will be particularly useful for our purpose.

Theorem 5.2 (Choi, Kim [6]). *The extreme points of the unit ball of $\mathcal{P}(^2\ell_\infty^2)$ are the polynomials of the form*

$$\pm x^2,\ \pm y^2,\ \pm(tx^2-ty^2\pm2\sqrt{t(1-t)}xy),$$

with $t\in[1/2,1]$.

Theorem 5.3 (Gámez-Merino, Muñoz-Fernández, Sánchez, Seoane-Sepúlveda [20]). *If $\mathcal{P}(^2\square)$ denotes the space $\mathcal{P}(^2\mathbb{R}^2)$ endowed with the sup norm over the unit interval $\square=[0,1]^2$ and $\mathsf{B}_\square$ is its unit ball, then the extreme points of $\mathsf{B}_\square$ are*

$$\pm(tx^2-y^2+2\sqrt{1-t}xy)\quad\textit{and}\quad\pm\left(-x^2+ty^2+2\sqrt{1-t}xy\right)\quad\textit{with } t\in[0,1]$$

or

$$\pm(x^2+y^2-xy),\ \pm(x^2+y^2-3xy),\ \pm x^2,\ \pm y^2.$$

Note that Theorem 5.3 is a kind on non-symmetric version of Theorem 5.2 and will be specially important when we are estimating the constants for $m \geq 4$. We refer the interested reader to [27,29] for other recent work on geometry of spaces of polynomials.

6. ESTIMATES FOR $D_{\mathbb{R},m}$

First observe that $\mathcal{P}(^m\ell_\infty^n)$ is a real linear space of dimension $N=\binom{m+n-1}{n-1}$. Therefore $\mathcal{P}(^m\ell_\infty^n)$ is algebraically isomorphic to $\mathbb{R}^N$ and a natural isomorphism between $\mathcal{P}(^m\ell_\infty^n)$ and $\mathbb{R}^N$ is given by $\mathbb{R}^N \ni \mathbf{a} \mapsto P_\mathbf{a} \in \mathcal{P}(^m\ell_\infty^n)$ where $P_\mathbf{a}(\mathbf{x}) := \sum_{|\alpha|=m} a_\alpha \mathbf{x}^\alpha$ is the m-homogeneous polynomial on $\mathbb{R}^n$ whose coefficients are the coordinates of the vector $\mathbf{a}$ assigned to each monomial $\mathbf{x}^\alpha$ in a prescribed order. Notice too that in order to caculate the optimal polynomial Bohnenblust-Hille constant for polynomials in $\mathcal{P}(\ell_\infty^n)$, which we denote by $D_{\mathbb{R},m}(\ell_\infty^n)$, it suffices to consider normalized polynomials. Then

$$D_{\mathbb{R},m}(\ell_\infty^n) := \sup\left\{\frac{\|\mathbf{a}\|_{\frac{2m}{m+1}}}{\|P_\mathbf{a}\|} : P_\mathbf{a} \in \mathcal{P}(^m\ell_\infty^n)\right\} = \sup\left\{\|\mathbf{a}\|_{\frac{2m}{m+1}} : P_\mathbf{a} \in \mathsf{B}_{\mathcal{P}(^m\ell_\infty^n)}\right\}, \tag{6.1}$$

where $\|\cdot\|_p$ denotes the ℓ_p norm. By convexity of $\|\cdot\|_p$ (notice that here $p=\frac{2m}{m+1}\geq 1$ for all $m\in\mathbb{N}$) we also have

$$D_{\mathbb{R},m}(\ell_\infty^n) = \sup\left\{\|\mathbf{a}\|_{\frac{2m}{m+1}} : P_\mathbf{a} \in \operatorname{ext}(\mathsf{B}_{\mathcal{P}(^m\ell_\infty^n)})\right\}. \tag{6.2}$$

6.1. **The case** $m=3$**.** Let us define $P_3 : \ell_\infty^6 \to \mathbb{R}$ by

$$P_3(x) = (x_1+x_2)\left(x_3^2+x_3x_4-x_4^2\right)+(x_1-x_2)\left(x_5^2+x_5x_6-x_6^2\right).$$

We have $\|P_3\| = 2\times\frac{5}{4}$. Also

$$\left(\sum_{|\alpha|=3}|a_\alpha|^{\frac{6}{4}}\right)^{\frac{4}{6}} \leq D_{\mathbb{R},3}\,\|P_3\|.$$

We, thus, have:

Proposition 6.1.

$$D_{\mathbb{R},3} \geq \frac{(4\times 3)^{4/6}}{2\times\frac{5}{4}} \approx 2.096.$$

6.2. **The case** $m=4$**.** It is not difficult to prove that the maximum value of $\frac{\|\mathbf{a}\|_{\frac{8}{5}}}{\|P_\mathbf{a}\|}$ where $P_\mathbf{a}$ ranges over the subspace of $\mathcal{P}(^4\ell_\infty^2)$ given by

$$\{ax^4+by^4+cx^2y^2 : a,b,c\in\mathbb{R}\},$$

is attained for the polynomial $Q_4(x,y)=x^4+y^4-3x^2y^2$:

Theorem 6.2. *If $E=\{ax^4+by^4+cx^2y^2 : a,b,c\in\mathbb{R}\}$ is endowed with the sup norm over the unit ball of ℓ_∞^2, then the optimal polynomial Bohnenblust-Hille constant for polynomials in E, which we denote by $D_{\mathbb{R},4}(E)$, is*

$$D_{\mathbb{R},4}(E) = \|(1,1,-3)\|_{\frac{8}{5}} = \left(2+(3)^{8/5}\right)^{5/8} \approx 3.610.$$

In particular

$$D_{\mathbb{R},4} \geq D_{\mathbb{R},4}(\ell_\infty^2) \geq D_{\mathbb{R},4}(E) \approx 3.610.$$

Moreover, equality is attained in the polynomial Bohnenblust-Hille inequality in E *for the polynomials* $P(x,y) = \pm(x^4 + y^4 - 3xy)$.

Proof. The same argument that led us to (6.2) also shows that

$$D_{\mathbb{R},4}(E) = \sup\left\{ \|\mathbf{a}\|_{\frac{8}{5}} : P_{\mathbf{a}} \in \mathrm{ext}(\mathsf{B}_E) \right\},$$

where $\mathrm{ext}(\mathsf{B}_E)$ is the set of extreme points in the unit ball, B_E, of E endowed with the supremum norm over the unit ball of ℓ_∞^2. Since $\mathrm{ext}(\mathsf{B}_E)$ is described in Theorem 5.3, we conclude that

$$\begin{aligned} D_{\mathbb{R},4}(E) &= \max\{\|(t,-1,2\sqrt{1-t})\|_{\frac{8}{5}}, \|(-1,t,2\sqrt{1-t})\|_{\frac{8}{5}}, \|(1,1,-1)\|_{\frac{8}{5}}, \|(1,1,-3)\|_{\frac{8}{5}} : \ t \in [0,1]\} \\ &= \|(1,1,-3)\|_{\frac{8}{5}} = \left(2 + (3)^{8/5}\right)^{5/8}. \end{aligned}$$

□

6.3. **Higher values of m.** We consider the polynomials

$$Q_{4k}(x,y) = \left(x^4 + y^4 - 3x^2y^2\right)^k,$$

for all $k \in \mathbb{N}$. Notice that $\|Q_{4k}\| = 1$ for all $k \in \mathbb{N}$. Therefore, using (6.1) together with the formula for the coefficients of the Q_{4k} given by (9.7), we can obtain estimates for $D_{\mathbb{R},4k}$ with k arbitrary (see Table 1). In fact we have:

Theorem 6.3. *If* $k \in \mathbb{N}$ *then*

$$D_{\mathbb{R},4k} \geq \left[\sum_{j=0}^{2k} |B_j|^{\frac{8k}{4k+1}} \right]^{\frac{4k+1}{8k}},$$

where

$$B_j = \sum_{\ell=0}^{\lfloor \frac{j}{2} \rfloor} \frac{k!(-3)^{j-2\ell}}{\ell!(j-2\ell)!(k-j+\ell)!},$$

for $j = 0, \ldots, 2k$.

$m = 8$	$D_{\mathbb{R},8} \geq 14.86998167$	$m = 80$	$D_{\mathbb{R},80} \geq 3.0496 \times 10^{13}$
$m = 12$	$D_{\mathbb{R},12} \geq 66.39260961$	$m = 120$	$D_{\mathbb{R},120} \geq 2.6821 \times 10^{20}$
$m = 16$	$D_{\mathbb{R},16} \geq 306.6665737$	$m = 160$	$D_{\mathbb{R},160} \geq 2.4320 \times 10^{27}$
$m = 20$	$D_{\mathbb{R},20} \geq 1442.799763$	$m = 200$	$D_{\mathbb{R},200} \geq 2.2443 \times 10^{34}$
$m = 24$	$D_{\mathbb{R},24} \geq 6866.770014$	$m = 240$	$D_{\mathbb{R},240} \geq 2.0924 \times 10^{41}$
$m = 28$	$D_{\mathbb{R},28} \geq 32940.16505$	$m = 280$	$D_{\mathbb{R},280} \geq 1.9649 \times 10^{48}$
$m = 32$	$D_{\mathbb{R},32} \geq 1.5892 \times 10^5$	$m = 320$	$D_{\mathbb{R},320} \geq 1.8549 \times 10^{55}$
$m = 36$	$D_{\mathbb{R},36} \geq 7.7009 \times 10^5$	$m = 360$	$D_{\mathbb{R},360} \geq 1.7582 \times 10^{62}$
$m = 40$	$D_{\mathbb{R},40} \geq 3.7444 \times 10^6$	$m = 400$	$D_{\mathbb{R},400} \geq 1.6718 \times 10^{69}$

TABLE 1. Estimates for $D_{\mathbb{R},m}$ for some values of m.

We also have the following representation, in the form C^m, of some of these lower bounds:

m	$D_{\mathbb{R},m} \geq$
200	$(1.48509930)^{200}$
800	$(1.49212548)^{800}$
$1,600$	$(1.49357368)^{1,600}$
$4,000$	$(1.49455267)^{4,000}$
$7,200$	$(1.49487590)^{7,200}$
$8,800$	$(1.49495333)^{8,800}$
$12,000$	$(1.49504910)^{12,000}$

6.4. **Comparing the optimal real and complex polynomial Bohnenblust–Hille constants.** Let us denote the optimal constants satisfying the polynomial Bohnenblust–Hille inequality, for degree m and scalar field $\mathbb{K}$, by $O_{\mathbb{K},m}$. From [9, 33], the best known estimates for the complex case are:

$$O_{\mathbb{C},m} \leq \begin{cases} \left(\frac{2}{\sqrt{\pi}}\right)^{m-1} \left(\frac{m^{\frac{m}{2}}(m+1)^{\frac{m+1}{2}}}{2^m (m!)^{\frac{m+1}{2m}}}\right) & \text{for } m = 2,3,4, \text{and} \\ \left(1+\frac{1}{m-1}\right)^{m-1} \sqrt{m}\left(\sqrt{2}\right)^{m-1} & \text{for } m \geq 5. \end{cases} \tag{6.3}$$

Thus, given $\varepsilon > 0$, there is a positive integer N_0 such that

$$O_{\mathbb{C},m} \leq \left(\sqrt{2}+\varepsilon\right)^m$$

whenever $m \geq N_0$. For $\varepsilon = 0.5$, for example, it can be checked that

$$D_{\mathbb{C},m} \leq (1.464)^m$$

whenever $m \geq 100$.

Hence, by using the estimates from the previous table we conclude that for $m = 4k$ and

$$200 \leq m \leq 1.2 \cdot 10^4$$

we have

$$O_{\mathbb{C},m} < O_{\mathbb{R},m}. \tag{6.4}$$

Let us stress that there is very little information on the comparison between $O_{\mathbb{C},m}$ and $O_{\mathbb{R},m}$ in the literature. For example, for $m = 3, 4$, using (6.3), Proposition 6.1 and Theorem 6.2, we have

$$O_{\mathbb{C},3} \leq 4.008 \text{ and } O_{\mathbb{R},3} \geq 2.086$$
$$O_{\mathbb{C},4} \leq 11.019 \text{ and } O_{\mathbb{R},4} \geq 3.616,$$

and hence (6.4) is unknown for these values of m.

The comparison between the optimal constants for the real and complex case in the multilinear case is also unknown. All that is known for multilinear mappings is that for 2-linear forms the optimal constant of the real case is strictly bigger that the constant of the complex case.

If the numerical computations from the previous table hold (following its trend) for arbitrarily large values of m then we will probably have (6.4) for all m, and we conjecture that this is actually what happens.

6.5. **On the optimal hypercontractivity constants.** Let $H_{\mathbb{R}}$ and $H_{\mathbb{C}}$ represent the optimal hypercontractivity constants, i.e.,

$$H_{\mathbb{K}} := \inf\{H > 0 : O_{\mathbb{K},m} \leq H^m \text{ for } m \text{ large enough}\}.$$

Using Theorem 3.1 together with (6.3) it follows immediately that

$$O_{\mathbb{R},m} \leq \begin{cases} \left(\frac{4}{\sqrt{\pi}}\right)^{m-1} \left(\frac{m^{\frac{m}{2}}(m+1)^{\frac{m+1}{2}}}{2^m (m!)^{\frac{m+1}{2m}}}\right) & \text{for } m = 2,3,4, \text{ and} \\ \left(1+\frac{1}{m-1}\right)^{m-1} \sqrt{m}\left(2\sqrt{2}\right)^{m-1} & \text{for } m \geq 5. \end{cases} \tag{6.5}$$

Now having in mind that $\left(1+\frac{1}{k}\right)^k < e$ for all $k \in \mathbb{N}$ and $\frac{e\sqrt{k}}{2\sqrt{2}} < b^k$ for all $k \geq 5$ with $b = \sqrt[5]{\frac{e\sqrt{5}}{2\sqrt{2}}}$, it is easy to prove that

$$O_{\mathbb{R},m} \leq (2\sqrt{2}b)^m,$$

for all $m \in \mathbb{N}$. Observe that $H_{\mathbb{R}} \leq 2\sqrt{2}b < 3.296$. By Theorem 4.1 we have

$$1.177 < H_{\mathbb{R}} < 3.296$$

and combining this result with the estimates from Theorem 6.3 we have

$$1.495 < H_{\mathbb{R}} < 3.296.$$

Actually, since the growth of $\left(1+\frac{1}{m-1}\right)^{m-1} \sqrt{m}\left(2\sqrt{2}\right)^{m-1}$ is dominated by $(2\sqrt{2}+\varepsilon)^m$ for any $\varepsilon > 0$, the following theorem summarizes our results:

Theorem 6.4. *The real hypercontractivity constant* $H_{\mathbb{R}}$ *is such that*

$$1.495 \leq H_{\mathbb{R}} \leq 2\sqrt{2} < 2.829.$$

The following problem still remains open.

Problem 6.5. *Is the hypercontractivity of the complex polynomial Bohnenblust–Hille inequality an optimal result?*

7. A LOWER ESTIMATE FOR $D_{\mathbb{C},2}$

Let $P : \ell_\infty^2(\mathbb{C}) \to \mathbb{C}$ be a 2-homogeneous polynomial given by

$$P(z_1, z_2) = az_1^2 + bz_1z_2 + cz_2^2,$$

with $a, b, c \in \mathbb{R}$. By the Maximum Modulus Principle we have

$$\begin{aligned} \|P\| &= \sup\{|az_1^2 + bz_1z_2 + cz_2^2| : |z_1| \leq 1,\ |z_2| \leq 1\} \\ &= \sup\{|az_1^2 + bz_1z_2 + cz_2^2| : |z_1| = |z_2| = 1\} \\ &= \sup\{|ae^{i2\theta_1} + be^{i\theta_1}e^{i\theta_2} + ce^{i\theta_2}| : \theta_1, \theta_2 \in [0, 2\pi]\} \\ &= \sup\{|ae^{i2(\theta_1-\theta_2)} + be^{i(\theta_1-\theta_2)} + c| : \theta_1 - \theta_2 \in [0, 2\pi]\} \\ &= \sup\{|az^2 + bz + c| : |z| = 1\} \\ &= \sup\{|az^2 + bz + c| : |z| \leq 1\}. \end{aligned}$$

In other words, $\|P\| = \|(a, b, c)\|_{\mathbb{C}}$, where

$$\|(a, b, c)\|_{\mathbb{C}} := \sup\{|az^2 + bz + c| : |z| \leq 1\}$$

is a norm on $\mathbb{R}^3$ studied in depth by Aron and Klimek in [2]. Using Aron and Klimek's results on the geometry of $\mathbb{R}^3$ with the norm $\|\cdot\|_{\mathbb{C}}$ we can prove the following:

Proposition 7.1. *Let $E_{\mathbb{R}}$ be the real subspace of $\mathcal{P}(^2\ell_\infty^2(\mathbb{C}))$ given by $\{az_1^2+bz_1z_2+cz_2^2 : (a,b,c)\in\mathbb{R}^3\}$. Then the optimal complex polynomial Bohnenblust-Hille constant for polynomials in $E_{\mathbb{R}}$, which we denote by $D_{\mathbb{C},2}$, is given by $D_{\mathbb{C},2}(E_{\mathbb{R}})=\sqrt[4]{\frac{3}{2}}$. In particular $D_{\mathbb{C},2}\geq\sqrt[4]{\frac{3}{2}}\approx 1.1066$.*

Proof. Using (6.2) for the subset $E_{\mathbb{R}}$ of $\ell_\infty^2(\mathbb{C})$ we have

$$\begin{aligned} D_{\mathbb{C},2}(E_{\mathbb{R}}) &= \sup\{\|(a,b,c)\|_{\frac{4}{3}} : \|az_1^2+bz_1z_2+cz_2^2\|\leq 1\} \\ &= \sup\{\|(a,b,c)\|_{\frac{4}{3}} : \|(a,b,c)\|_{\mathbb{C}}\leq 1\} \\ &= \sup\{\|(a,b,c)\|_{\frac{4}{3}} : (a,b,c)\in \operatorname{ext}(\mathsf{B}_{E_{\mathbb{R}}})\}, \end{aligned}$$

where $\operatorname{ext}(\mathsf{B}_{E_{\mathbb{R}}})$ is the set of extreme points of the unit ball of $E_{\mathbb{R}}$, namely $\mathsf{B}_{E_{\mathbb{R}}}$. The set $\operatorname{ext}(\mathsf{B}_{E_{\mathbb{R}}})$ has been explicitly described by Aron and Klimek [2] and turns out to be

$$\operatorname{ext}(\mathsf{B}_{E_{\mathbb{R}}})=\left\{\left(s,\sqrt{4|s||t|\left(\frac{1}{(|s|+|t|)^2}-1\right)},t\right) : (s,t)\in G\right\},$$

where $G=\{(s,t)\in\mathbb{R}^2 : |s|+|t|<1 \text{ and } |s+t|\leq (s+t)^2\}\cup\{\pm(1,0),\pm(0,1)\}$. Hence

$$D_{\mathbb{C},2}(E_{\mathbb{R}})=\sup\left\{\left(|s|^{\frac{4}{3}}+|t|^{\frac{4}{3}}\left[4|s||t|\left(\frac{1}{(|s|+|t|)^2}-1\right)\right]^{\frac{2}{3}}\right)^{\frac{3}{4}} : (s,t)\in G\right\}.$$

If $\Phi(s,t)=\left(|s|^{\frac{4}{3}}+|t|^{\frac{4}{3}}\left[4|s||t|\left(\frac{1}{(|s|+|t|)^2}-1\right)\right]^{\frac{2}{3}}\right)^{\frac{3}{4}}$ for $(s,t)\in G$, one can prove using elementary calculus that Φ attains its maximum on G at $\pm\left(\frac{\sqrt{3}}{6},-\frac{\sqrt{3}}{6}\right)$ and $\Phi\left(\frac{\sqrt{3}}{6},-\frac{\sqrt{3}}{6}\right)=\sqrt[4]{\frac{3}{2}}$.

The reader can find a sketch of the graph of Φ on the part of G contained in the second quadrant in Figure 1.

□

Combining the previous result and the known fact that $D_{\mathbb{C},2}\leq 1.7431$ we have the following result:

Theorem 7.2.

$$1.1066\leq D_{\mathbb{C},2}\leq 1.7431.$$

8. Multilinear variants of the Bohnenblust–Hille inequality

Although the work of Bohnenblust and Hille is focused on complex scalars, it is well-known that the result also holds for real scalars:

If A is a continuous n-linear form on $c_0\times\cdots\times c_0$, then there is a constant $C_{\mathbb{K},n}$ (depending only on n and $\mathbb{K}$) such that

$$\left(\sum_{i_1,\ldots,i_m=1}^{\infty}|A(e_{i_1},\ldots,e_{i_n})|^{\frac{2n}{n+1}}\right)^{\frac{n+1}{2n}}\leq C_{\mathbb{K},n}\|A\|.$$



Figure 1. Graph of Φ on G intersected with the second quadrant.

The estimates for $C_{\mathbb{K},n}$ were improved along the decades (see [7,21,33]). From recent works (see [18,32]) we know that, for real scalars,

$$\begin{aligned} C_{\mathbb{R},2} &= \sqrt{2} \approx 1.414 \\ 1.587 \leq C_{\mathbb{R},3} &\leq 1.782 \\ 1.681 \leq C_{\mathbb{R},4} &\leq 2 \\ 1.741 \leq C_{\mathbb{R},5} &\leq 2.298 \\ 1.811 \leq C_{\mathbb{R},6} &\leq 2.520 \end{aligned}$$

and, for the complex case,

$$\begin{aligned} C_{\mathbb{C},2} &\leq \left(\frac{2}{\sqrt{\pi}}\right) \approx 1.128 \\ C_{\mathbb{C},3} &\leq 1.273 \\ C_{\mathbb{C},4} &\leq 1.437 \\ C_{\mathbb{C},5} &\leq 1.621 \\ C_{\mathbb{C},10} &\leq 2.292 \\ C_{\mathbb{C},15} &\leq 2.805. \end{aligned}$$

The lower bounds for $C_{\mathbb{R},m}$ obtained in [18] are $2^{\frac{m-1}{m}}$, so the precise value for $C_{\mathbb{R},m}$ for "big m's" is quite uncertain. Very recently, it was shown that for both real and complex scalars the asymptotic behavior of the best values for $C_{\mathbb{K},n}$ is optimal [17].

The (complex and real) Bohnenblust–Hille inequality can be re-written in the context of multiple summing multilinear operators.

Let $X_1, \ldots, X_m$ and Y be Banach spaces over $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$, and X' be the topological dual of X. By $\mathcal{L}(X_1, \ldots, X_m; Y)$ we denote the Banach space of all continuous m-linear mappings from $X_1 \times \cdots \times X_m$ to Y with the usual sup norm. For $x_1, ..., x_n$ in X, let

$$\|(x_j)_{j=1}^n\|_{w,1} := \sup\{\|(\varphi(x_j))_{j=1}^n\|_1 : \varphi \in X', \|\varphi\| \leq 1\}.$$

If $1 \leq p < \infty$, an m-linear mapping $U \in \mathcal{L}(X_1, \ldots, X_m; Y)$ is multiple $(p;1)$-summing (denoted $\Pi_{(p;1)}(X_1, \ldots, X_m; Y)$) if there exists a constant $U_{\mathbb{K},m} \geq 0$ such that

$$\left(\sum_{j_1,\ldots,j_m=1}^{N} \left\| U(x_{j_1}^{(1)}, \ldots, x_{j_m}^{(m)}) \right\|^p \right)^{\frac{1}{p}} \leq U_{\mathbb{K},m} \prod_{k=1}^{m} \left\| (x_j^{(k)})_{j=1}^N \right\|_{w,1} \tag{8.1}$$

for every $N \in \mathbb{N}$ and any $x_{j_k}^{(k)} \in X_k$, $j_k = 1, \ldots, N$, $k = 1, \ldots, m$. The infimum of the constants satisfying (8.1) is denoted by $\|U\|_{\pi(p;1)}$. For $m = 1$ we recover the well-known concept of absolutely $(p;1)$-summing operators (see, e.g. [8, 16]).

The Bohnenblust–Hille inequality can be re-written in the context of multiple summing multilinear operators in the following sense: every continuous m-linear form $U : X_1 \times \cdots \times X_m \to \mathbb{K}$ is multiple $(\frac{2m}{m+1}; 1)$-summing. Moreover

$$\|U\|_{\pi(\frac{2m}{m+1};1)} \leq C_{\mathbb{K},m} \|U\|. \tag{8.2}$$

For details we refer to [14] and references therein.

From now on if $P : X \to Y$ is a m-homogeneous polynomial then $\overset{\vee}{P}$ denotes the (unique) symmetric m-linear map (also called the *polar* of P) associated to P. Recall that an m-homogeneous polynomial $P : X \to Y$ is multiple $(p;1)$-summing (denoted $\mathcal{P}_{(p;1)}(^m X; Y)$) if there exists a constant $P_{\mathbb{K},m} \geq 0$ such that

$$\left(\sum_{j_1,\ldots,j_m=1}^{N} \left\| \overset{\vee}{P}(x_{j_1}^{(1)}, \ldots, x_{j_m}^{(m)}) \right\|^p \right)^{\frac{1}{p}} \leq P_{\mathbb{K},m} \prod_{k=1}^{m} \left\| (x_j^{(k)})_{j=1}^N \right\|_{w,1} \tag{8.3}$$

for every $N \in \mathbb{N}$ and any $x_{j_k}^{(k)} \in X$, $j_k = 1, \ldots, N$, $k = 1, \ldots, m$. The infimum of the constants satisfying (8.3) is denoted by $\|P\|_{\pi(p;1)}$. Note that

$$\|P\|_{\pi(p;1)} = \left\| \overset{\vee}{P} \right\|_{\pi(p;1)}.$$

If $P \in \mathcal{P}(^m X; \mathbb{K})$ then $\overset{\vee}{P} \in \mathcal{L}(^m X; \mathbb{K}) = \Pi_{(\frac{2m}{m+1};1)}(^m X; \mathbb{K})$ and

$$\|P\|_{\pi(\frac{2m}{m+1};1)} = \left\| \overset{\vee}{P} \right\|_{\pi(\frac{2m}{m+1};1)} \overset{(8.2)}{\leq} C_{\mathbb{K},m} \left\| \overset{\vee}{P} \right\| \leq \frac{m^m}{m!} C_{\mathbb{K},m} \|P\|.$$

So, since $C_{\mathbb{K},m}$ does not depend on X and P we conclude that there are constants $L_{\mathbb{K},m}$ (which do not depend on X and P) such that

$$\left(\sum_{j_1,\ldots,j_m=1}^{N}\left\|\overset{\vee}{P}(x_{j_1}^{(1)},\ldots,x_{j_m}^{(m)})\right\|^{p}\right)^{\frac{1}{p}}\leq L_{\mathbb{K},m}\left\|P\right\|\prod_{k=1}^{m}\left\|(x_j^{(k)})_{j=1}^{N}\right\|_{w,1}.$$

Note that if $X=\ell_\infty^N$, and $x^{(j)}=e_j$ for every $j=1,...,N$, since

$$\left\|(x^{(j)})_{j=1}^{N}\right\|_{w,1}=1,$$

we have

$$\left(\sum_{j_1,\ldots,j_m=1}^{N}\left\|\overset{\vee}{P}(e_{j_1},\ldots,e_{j_m})\right\|^{\frac{2m}{m+1}}\right)^{\frac{m+1}{2m}}\leq L_{\mathbb{K},m}\left\|P\right\| \tag{8.4}$$

for every $N\in\mathbb{N}$, which can be regarded as a “kind of” polynomial Bohnenblust–Hille inequality. Since (8.4) is confined to the symmetric case, there is no obvious relation between the optimal values for $C_{\mathbb{K},m}$ and the optimal values of $L_{\mathbb{K},m}$.

For $m=2$ it is well-known that $C_{\mathbb{R},2}=\sqrt{2}$. For $m>2$ the precise values of $C_{\mathbb{R},m}$ are not known. Since

$$L_{\mathbb{R},m}\leq\frac{m^m}{m!}C_{\mathbb{R},m},$$

we have

$$\begin{aligned}L_{\mathbb{R},2}&\leq 2.828\\ L_{\mathbb{R},3}&\leq 8.018\\ L_{\mathbb{R},4}&\leq 21.333\end{aligned}$$

In the following section we shall provide nontrivial lower bounds for $L_{\mathbb{R},m}$.

9. ESTIMATES FOR $L_{\mathbb{R},m}$

Definition 9.1. *Let $d=\binom{m+n-1}{n-1}$ be the dimension of the space of all m-homogeneous polynomials on $\mathbb{R}^n$. For every $m,n\in\mathbb{N}$, we define $\Phi_{m,n}:\mathbb{R}^d\to\mathbb{R}$ as follows: Take $\mathbf{a}\in\mathbb{R}^d$ and consider the the m-homogeneous polynomial $P_{\mathbf{a}}(\mathbf{x})=\sum_{|\alpha|=m}a_\alpha\mathbf{x}^\alpha$ whose coefficients are the coordinates of $\mathbf{a}$. In order to avoid redundancies, assume that $\mathbf{a}=(a_\alpha)$ where the coordinates are arranged according to the lexicographic order of the α's. Then if $L_{\mathbf{a}}$ is the polar of $P_{\mathbf{a}}$ we define*

$$\Phi_{m,n}(\mathbf{a}):=\left[\sum_{i_1+\cdots+i_m=m}\left|L_{\mathbf{a}}(e_{i_1},\ldots,e_{i_m})\right|^{\frac{2m}{m+1}}\right]^{\frac{m+1}{2m}}.$$

Notice that Lemma 2.1 allows us to write $\Phi_{m,n}$ as

$$\begin{aligned}\Phi_{m,n}(\mathbf{a}) &= \left[\sum_{\substack{\alpha=(\alpha_1,\cdots,\alpha_n)\\|\alpha|=m}} \binom{m}{\alpha} |L_{\mathbf{a}}(e_1^{\alpha_1},\ldots,e_n^{\alpha_n})|^{\frac{2m}{m+1}}\right]^{\frac{m+1}{2m}} \\ &= \left[\sum_{|\alpha|=m}\binom{m}{\alpha}\left|\frac{a_\alpha}{\binom{m}{\alpha}}\right|^{\frac{2m}{m+1}}\right]^{\frac{m+1}{2m}}. \qquad (9.1)\end{aligned}$$

Also $\Phi_{m,n}$ is, essentially, the composition of the norm in $\ell^d_{\frac{2m}{m+1}}$ with the natural isomorphism between $\mathcal{L}^s({}^m\mathbb{R}^n)$ and $\mathcal{P}({}^m\mathbb{R}^n)$. Therefore $\Phi_{m,n}$ is convex and by virtue of Krein–Milman Theorem

$$L_{\mathbb{R},m} \geq L_{\mathbb{R},m}\left(\ell_\infty^n\right) := \sup\{\Phi_{m,n}(\mathbf{a}) : \mathbf{a} \in \mathsf{B}_{\mathcal{P}({}^m\ell_\infty^n)}\} = \sup\{\Phi_{m,n}(\mathbf{a}) : \mathbf{a} \in \operatorname{ext}(\mathsf{B}_{\mathcal{P}({}^m\ell_\infty^n)})\},$$

where $\operatorname{ext}(\mathsf{B}_{\mathcal{P}({}^m\ell_\infty^n)})$ is the set of extreme points of $\mathsf{B}_{\mathcal{P}({}^m\ell_\infty^n)}$ and $L_{\mathbb{R},m}\left(\ell_\infty^n\right)$ is the optimal constant in the Bohnenblust-Hille type inequality (8.4) for real polynomials in $\mathcal{P}({}^m\ell_\infty^n)$. Observe that even in the case where the geometry of $\mathsf{B}_{\mathcal{P}({}^m\ell_\infty^n)}$ is not know, the mapping $\Phi_{m,n}$ provides a lower bound for $L_{\mathbb{R},m}$, namely

$$L_{\mathbb{R},m} \geq \frac{\Phi_{m,n}(\mathbf{a})}{\|P_{\mathbf{a}}\|}, \qquad (9.2)$$

for all $\mathbf{a} \in \mathbb{R}^d$.

In the following we will try to use the fact that the extreme points of $\mathsf{B}_{\mathcal{P}({}^m\ell_\infty^n)}$ have been characterized for some choices of m and n (see for instance Theorem 5.2).

9.1. **Case** $m = 2$**.** We begin by illustrating that even sharp information for lower estimates for $C_{\mathbb{R},2}$ may be useless for evaluating lower estimates for $L_{\mathbb{R},2}$. For instance, if $m = 2$ in the multilinear Bohnenblust–Hille inequality (in fact, Littlewood's 4/3 inequality, [24]) the best constant is $C_{\mathbb{R},2} = \sqrt{2}$ and this estimate is achieved (see [18]) when we use the bilinear form $T_2 : \ell_\infty^2 \times \ell_\infty^2 \to \mathbb{R}$ given by

$$T_2(x,y) = x_1y_1 + x_1y_2 + x_2y_1 - x_2y_2.$$

Note that T_2 is symmetric and the polynomial associated to T_2 is $P_2 : \ell_\infty^2 \to \mathbb{R}$ given by

$$P_2(x) = x_1^2 + 2x_1x_2 - x_2^2.$$

Since $\|P_2\| = \|T_2\| = 2$, the constant $L_{\mathbb{R},2}$ that appears for this choice of P_2 is again $\sqrt{2}$, which is far from being a good lower estimate, as we shall see in the next result, that gives the exact value for the constant $L_{\mathbb{R},2}(\ell_\infty^2)$.

Theorem 9.2. $L_{\mathbb{R},2} \geq 1.7700$. *More precisely,*

$$L_{\mathbb{R},2}\left(\ell_\infty^2\right) = \sup\left\{\left[2t^{\frac{4}{3}} + 2\left(\sqrt{t(1-t)}\right)^{\frac{4}{3}}\right]^{\frac{3}{4}} : t \in [1/2,1]\right\} \approx 1.7700$$

and the supremum is attained at $t_0 \approx 0.9147$.

Proof. Observe that for polynomials in $\mathcal{P}(^2\ell_\infty^2)$ of the form $P_{\mathbf{a}}(x,y) = ax^2 + by^2 + cxy$ with $\mathbf{a} = (a,b,c)$ we have

$$\Phi_{2,2}(a,b,c) = \left[a^{\frac{4}{3}} + b^{\frac{4}{3}} + 2\left(\frac{c}{2}\right)^{\frac{4}{3}}\right]^{\frac{3}{4}}. \tag{9.3}$$

Using the Krein–Milman approach

$$L_{\mathbb{R},2}\left(\ell_\infty^2\right) = \sup\{\Phi_{22}(\mathbf{a}) : \mathbf{a} \in \operatorname{ext}(\mathsf{B}_{\mathcal{P}(^2\ell_\infty^2)})\}.$$

Now, by Theorem 5.2, $\operatorname{ext}(\mathsf{B}_{\mathcal{P}(^2\ell_\infty^2)})$ consists of the polynomials

$$\pm(1,0,0), \quad \pm(0,1,0) \quad \text{and} \quad \pm\left(t, -t, \pm 2\sqrt{t(1-t)}\right),$$

with $t \in [1/2, 1]$. Since the contribution of $\pm(1,0,0)$ and $\pm(0,1,0)$ to the supremum is irrelevant, we end up with

$$\begin{aligned} L_{\mathbb{R},2}\left(\ell_\infty^2\right) &= \sup\{\Phi_{2,2}(\pm(t,-t,\pm 2\sqrt{t(1-t)})) : t \in [1/2,1]\} \\ &= \sup\left\{\left[2t^{\frac{4}{3}} + 2\left(\sqrt{t(1-t)}\right)^{\frac{4}{3}}\right]^{\frac{3}{4}} : t \in [1/2,1]\right\}. \end{aligned}$$

The problem of maximizing (explicitly) this function is quite hard and the final result is far from being "good looking". The interested reader can obtain an explicit solution in radical form using a variety of symbolic computational software. A 4-digit approximation yields

$$L_{\mathbb{R},2} \geq L_{\mathbb{R},2}\left(\ell_\infty^2\right) \approx 1.7700,$$

where the maximum is attained at $t_0 \approx 0.9147$. $\square$

Remark 9.3. *A very good approximation of $L_{\mathbb{R},2}\left(\ell_\infty^2\right)$ can be obtained considering the polynomial*

$$P_{\mathbf{a}}(x,y) = x^2 - y^2 + xy,$$

i.e., $\mathbf{a} = (1,-1,1)$*. It is easy to check that $\|P_{\mathbf{a}}\| = 5/4$. Hence, using* (9.2) *we have*

$$L_{\mathbb{R},2}\left(\ell_\infty^2\right) \geq \frac{\Phi_{2,2}(1,-1,1)}{\|P_{\mathbf{a}}\|} = \frac{4}{5} \cdot \left(2 + 2\left(\frac{1}{2}\right)^{4/3}\right)^{3/4} \approx 1.728.$$

9.2. **Case** $m = 4$**.** In this section we calculate the exact value of $L_{\mathbb{R},4}$ in a subspace of $\mathcal{L}^s(^4\ell_\infty^2)$. Observe that the value of $L_{\mathbb{R},4}$ in a subspace is, obviously, a lower bound for $L_{\mathbb{R},4}$.

Theorem 9.4. *If $E = \{ax^4 + by^4 + cx^2y^2 : a, b, c \in \mathbb{R}\}$ and $\overset{\vee}{E}$ is the space of polars of elements in E endowed with the sup norm over the unit ball of ℓ_∞^2, then the optimal constant in* (8.4) *for elements in $\overset{\vee}{E}$ is given by*

$$L_{\mathbb{R},4}(\overset{\vee}{E}) = \left[2 + 6\left(\frac{1}{2}\right)^{\frac{8}{5}}\right]^{\frac{5}{8}} \approx 2.371.$$

In particular

$$L_{\mathbb{R},4} \geq L_{\mathbb{R},4}(\ell_\infty^2) \geq L_{\mathbb{R},4}(\overset{\vee}{E}) \approx 2.371.$$

Moreover, equality is attained in the Bohnenblust-Hille inequality in $\overset{\vee}{E}$ for the polars of the polynomials $P(x,y) = \pm(x^4 - y^4 + 3xy)$.

Proof. We just need to calculate the maximum of $\Phi_{4,2}$ over E, which is trivially isometric to the space $\mathcal{P}(^2\Box)$ (see Theorem 5.3 for the definition of $\mathcal{P}(^2\Box)$). If $\Phi = \Phi_{4,2}|_{\mathcal{P}(^2\Box)}$, then Φ is obviously convex and we have

$$\begin{aligned} L_{\mathbb{R},4} \geq L_{\mathbb{R},4}\left(\ell_\infty^2\right) &= \sup\{\Phi_{4,2}(\mathbf{a}) : \mathbf{a} \in \mathsf{B}_{\mathcal{P}(^4\ell_\infty^2)}\} \\ &\geq \sup\{\Phi(\mathbf{a}) : \mathbf{a} \in \mathsf{B}_{\mathcal{P}(^2\Box)}\} \\ &= \sup\{\Phi(\mathbf{a}) : \mathbf{a} \in \operatorname{ext}(\mathsf{B}_{\mathcal{P}(^2\Box)})\}, \end{aligned}$$

where the last equality is due to the Krein–Milman Theorem. Now by (9.1) we have

$$\Phi(a,b,c) = \left[a^{\frac{8}{5}} + b^{\frac{8}{5}} + 6\left(\frac{c}{6}\right)^{\frac{8}{5}}\right]^{\frac{5}{8}}.$$

Using Theorem 5.3 we obtain

$$\begin{aligned} &\sup\{\Phi(\mathbf{a}) : \mathbf{a} \in \mathsf{B}_{\mathcal{P}(^2\Box)}\} = \\ &= \max\left\{\left[1 + t^{\frac{8}{5}} + 6\left(\frac{\sqrt{1-t}}{3}\right)^{\frac{8}{5}}\right]^{\frac{5}{8}}, \left[2 + 6\left(\frac{1}{6}\right)^{\frac{8}{5}}\right]^{\frac{5}{8}}, \left[2 + 6\left(\frac{1}{2}\right)^{\frac{8}{5}}\right]^{\frac{5}{8}} : t \in [0,1]\right\} \\ &= \left[2 + 6\left(\frac{1}{2}\right)^{\frac{8}{5}}\right]^{\frac{5}{8}}. \end{aligned}$$

Observe that the maximum is attained at the polynomials $P(x,y) = \pm(x^4 + y^4 - 3xy)$. Hence we have proved that

$$L_{\mathbb{R},4} \geq \left[2 + 6\left(\frac{1}{2}\right)^{\frac{8}{5}}\right]^{\frac{5}{8}} \approx 2.371,$$

moreover, a better (bigger) lower estimate for $L_{\mathbb{R},4}$ cannot be obtained by considering polynomials of the form $ax^4 + by^4 + cx^2y^2$ with $a, b, c \in \mathbb{R}$. □

9.3. **Higher values of** m**.** The previous sections allow us to obtain lower estimates for $L_{\mathbb{R},m}$ for arbitrary large m's. In this section we consider polynomials of the form $P_{2k}(x,y) = (ax^2 + by^2 + cxy)^k$. In the following, if $h \in \mathbb{Z}$, $\lfloor h \rfloor$ denotes the biggest integer H so that $H \leq h$.

Proposition 9.5. *If $P_{2k}(x,y) = (ax^2 + by^2 + cxy)^k$, then $P_{2k}(x,y) = \sum_{j=0}^{2k} A_j x^j y^{2k-j}$ with*

$$A_j = \sum_{\ell=0}^{\lfloor \frac{j}{2} \rfloor} \frac{k! a^\ell b^{k-j+\ell} c^{j-2\ell}}{\ell!(j-2\ell)!(k-j+\ell)!}, \tag{9.4}$$

for $j = 0, \ldots, 2k$.

Proof. Using the multinomial formula:

$$P_{2k}(x,y) = (ax^2 + by^2 + cxy)^k = \sum_{\substack{\alpha_1+\alpha_2+\alpha_3=k \\ \alpha_1,\alpha_2,\alpha_3 \geq 0}} \frac{k!}{\alpha_1!\alpha_2!\alpha_3!} a^{\alpha_1} b^{\alpha_2} c^{\alpha_3} x^{2\alpha_1+\alpha_3} y^{2\alpha_2+\alpha_3}.$$

Therefore, $x^j y^{2n-j} = x^{2\alpha_1+\alpha_3} y^{2\alpha_2+\alpha_3}$ for $j = 1, \ldots, 2k$ implies that

$$\begin{cases} 2\alpha_1 + \alpha_3 = j, \\ 2\alpha_2 + \alpha_3 = 2k - j, \end{cases}$$

which, together with the fact that $\alpha_1+\alpha_2+\alpha_3=k$ and $\alpha_1,\alpha_2,\alpha_3\geq 0$ yield

$$\begin{cases} \alpha_3 = j-2\alpha_1, \\ \alpha_2 = k-j+\alpha_1, \end{cases}$$

with $\alpha_1=0,\dots,\lfloor \frac{j}{2}\rfloor$. As a result of the previous comments, the coefficient A_j is given by (9.4). □

Corollary 9.6. *If $k\in\mathbb{N}$ then*

$$L_{\mathbb{R},2k} \geq \left[\sum_{j=0}^{2k}\binom{2k}{j}\left|\frac{A_j}{\binom{2k}{j}}\right|^{\frac{4k}{2k+1}}\right]^{\frac{2k+1}{4k}}, \tag{9.5}$$

where

$$A_j=\sum_{\ell=0}^{\lfloor \frac{j}{2}\rfloor}\frac{k!(-1)^{k-j+\ell}t_0^{k-j+2\ell}(2\sqrt{t_0(1-t_0)})^{j-2\ell}}{\ell!(j-2\ell)!(k-j+\ell)!},$$

for $j=0,\dots,2k$ and t_0 is as in Theorem 9.2.

Proof. If $P_{2k}(x,y)=(ax^2+by^2+cxy)^k$, using (9.1), (9.2) and Proposition 9.5 we arrive at

$$L_{\mathbb{R},2k} \geq \frac{1}{\|P_{2k}\|}\left[\sum_{j=0}^{2k}\binom{2k}{j}\left|\frac{A_j}{\binom{2k}{j}}\right|^{\frac{4k}{2k+1}}\right]^{\frac{2k+1}{4k}},$$

with A_j as in (9.4). Then the corollary follows by considering the polynomial

$$P_{2k}(x,y)=(t_0x^2-t_0y^2+2\sqrt{t_0(1-t_0)}xy)^{2k},$$

which has norm 1. □

Hence (9.5) provides a systematic formula to obtain a lower bound for $L_{\mathbb{R},m}$ for even m's. Observe that for $k=2$ we have

$$L_{\mathbb{R},4} \geq \left[2t_0^{\frac{16}{5}}+6\left(\frac{2t_0-3t_0^2}{3}\right)^{\frac{8}{5}}+8(t_0\sqrt{t_0(1-t_0)})^{\frac{8}{5}}\right]^{\frac{5}{8}} \approx 2.1595,$$

which is a slightly worse constant than the one obtained in Section 9.2. Actually, the estimates (9.5) can be improved for multiples of 4. Indeed, we just need to consider the polynomials

$$Q_{4k}(x,y)=(ax^4+by^4+cx^2y^2)^k,$$

with $k\in\mathbb{N}$. Using exactly the same procedure described in this section

$$L_{\mathbb{R},4k} \geq \frac{1}{\|Q_{4k}\|}\left[\sum_{j=0}^{2k}\binom{4k}{2j}\left|\frac{A_j}{\binom{4k}{2j}}\right|^{\frac{8k}{4k+1}}\right]^{\frac{4k+1}{8k}},$$

where the A_j's, with $j=1,\dots,2k$ are the same as in (9.4). Now, putting $a=1$, $b=1$ and $c=-3$, i.e., considering powers of the extreme polynomial that appeared in Section 9.2, we would have that $\|Q_{4k}\|=1$ for all $k\in\mathbb{N}$, which proves the following:

Theorem 9.7. *If $k \in \mathbb{N}$ then*

$$L_{\mathbb{R},4k} \geq \left[\sum_{j=0}^{2k} \binom{4k}{2j} \left| \frac{B_j}{\binom{4k}{2j}} \right|^{\frac{8k}{4k+1}} \right]^{\frac{4k+1}{8k}}, \tag{9.6}$$

where

$$B_j = \sum_{\ell=0}^{\lfloor \frac{j}{2} \rfloor} \frac{k!(-3)^{j-2\ell}}{\ell!(j-2\ell)!(k-j+\ell)!}, \tag{9.7}$$

for $j = 0, \ldots, 2k$.

As an example, let us apply (9.6) and (9.7) to obtain estimates for $L_{\mathbb{R},8}$ and $L_{\mathbb{R},12}$. The polynomials are

$$\begin{aligned} Q_8(x,y) &= x^8 - 6x^6y^2 + 11x^4y^4 - 6x^2y^6 + y^8, \\ Q_{12}(x,y) &= x^{12} - 9x^{10}y^2 + 30x^8y^4 - 45x^6y^6 + 30x^4y^8 - 9x^2y^{10} + y^{12}. \end{aligned}$$

Then

$$L_{\mathbb{R},8} \geq \left[2 + 2\binom{8}{2} \left(\frac{6}{\binom{8}{2}} \right)^{\frac{16}{9}} + \binom{8}{4} \left(\frac{11}{\binom{8}{4}} \right)^{\frac{16}{9}} \right]^{\frac{9}{16}} \approx 3.2725.$$

$$L_{\mathbb{R},12} \geq \left[2 + 2\binom{12}{2} \left(\frac{9}{\binom{12}{2}} \right)^{\frac{24}{13}} + 2\binom{12}{4} \left(\frac{30}{\binom{12}{4}} \right)^{\frac{24}{13}} + \binom{12}{6} \left(\frac{45}{\binom{12}{6}} \right)^{\frac{24}{13}} \right]^{\frac{13}{24}} \approx 4.2441.$$

For higher degrees see Table 2.

$k=4$	$L_{\mathbb{R},16} \geq 5.390975019$	$k=40$	$L_{\mathbb{R},160} \geq 16805.46318$
$k=5$	$L_{\mathbb{R},20} \geq 6.787708182$	$k=50$	$L_{\mathbb{R},200} \geq 1.5654 \times 10^5$
$k=6$	$L_{\mathbb{R},24} \geq 8.511696468$	$k=60$	$L_{\mathbb{R},240} \geq 1.4581 \times 10^6$
$k=9$	$L_{\mathbb{R},36} \geq 16.65124974$	$k=90$	$L_{\mathbb{R},360} \geq 1.1781 \times 10^9$
$k=10$	$L_{\mathbb{R},40} \geq 20.81051033$	$k=100$	$L_{\mathbb{R},400} \geq 1.0972 \times 10^{10}$

TABLE 2. $L_{\mathbb{R},4k}$ for some values of $k \in \mathbb{N}$.

In order to clarify what the asymptotic growth of the sequence $(L_{\mathbb{R},4k})_{k\in\mathbb{N}}$ is, a simple calculation of the quotients of the estimates obtained in Table 2 for higher values of k indicates that the ratio of the estimates on $L_{\mathbb{R},4(k+1)}$ and $L_{\mathbb{R},4k}$ seem to tend to $\frac{5}{4}$.

As we have seen, in the real case we were able to deal with the case $m \geq 2$ even in the absence of information on the geometry of the unit ball of $\mathcal{P}(^m\ell_\infty^n)$. However, in the complex case the technique seemed less effective for $m \geq 3$. In order to obtain lower estimates for $D_{\mathbb{C},m}$, with $m \geq 3$ and sharper estimates for $D_{\mathbb{R},m}$, we believe that some effort should be made to obtain more information on the geometry of the unit ball of $\mathcal{P}(^m\ell_\infty^n)$ for higher values of m, n. We certainly hope that the present work may serve as a motivation for future works investigating the geometry of the unit ball of complex and real polynomial spaces on ℓ_∞^n.

References

[1] R. Aron, B. Beauzamy, and P. Enflo, *Polynomials in many variables: real vs complex norms*, J. Approx. Theory **74** (1993), no. 2, 181–198.

[2] R. M. Aron and M. Klimek, *Supremum norms for quadratic polynomials*, Arch. Math. (Basel) **76** (2001), no. 1, 73–80.

[3] H. P. Boas, *The football player and the infinite series*, Notices Amer. Math. Soc. **44** (1997), 1430–1435.

[4] H. P. Boas and D. Khavinson, *Bohr's power series theorem in several variables*, Proc. Amer. Math. Soc. **125** (1997), no. 10, 2975–2979.

[5] H. F. Bohnenblust and E. Hille, *On the absolute convergence of Dirichlet series*, Ann. of Math. (2) **32** (1931), no. 3, 600–622.

[6] Y. S. Choi and S. G. Kim, *The unit ball of* $\mathcal{P}({}^2l_2^2)$, Arch. Math. (Basel) **71** (1998), no. 6, 472–480.

[7] A. M. Davie, *Quotient algebras of uniform algebras*, J. London Math. Soc. (2) **7** (1973), 31–40.

[8] A. Defant and K. Floret, *Tensor norms and operator ideals*, North-Holland Mathematics Studies, vol. 176, North-Holland Publishing Co., Amsterdam, 1993.

[9] A. Defant, L. Frerick, J. Ortega-Cerdà, M. Ounaïes, and K. Seip, *The Bohnenblust–Hille inequality for homogeneous polynomials is hypercontractive*, Ann. of Math. (2) **174** (2011), no. 1, 485–497.

[10] A. Defant, J. C. Díaz, D. García, and M. Maestre, *Unconditional basis and Gordon-Lewis constants for spaces of polynomials*, J. Funct. Anal. **181** (2001), no. 1, 119–145.

[11] A. Defant, D. García, M. Maestre, and D. Pérez-García, *Bohr's strip for vector valued Dirichlet series*, Math. Ann. **342** (2008), no. 3, 533–555.

[12] A. Defant, D. García, M. Maestre, and P. Sevilla-Peris, *Bohr's strips for Dirichlet series in Banach spaces.* part 2, Funct. Approx. Comment. Math. **44** (2011), no. part 2, 165–189.

[13] A. Defant, M. Maestre, and U. Schwarting, *Bohr radii of vector valued holomorphic functions*, Adv. Math. **231** (2012), no. 5, 2837–2857.

[14] A. Defant, D. Popa, and U. Schwarting, *Coordinatewise multiple summing operators in Banach spaces*, J. Funct. Anal. **259** (2010), no. 1, 220–242.

[15] A. Defant and P. Sevilla-Peris, *A new multilinear insight on Littlewood's 4/3-inequality*, J. Funct. Anal. **256** (2009), no. 5, 1642–1664.

[16] J. Diestel, H. Jarchow, and A. Tonge, *Absolutely summing operators*, Cambridge Studies in Advanced Mathematics, vol. 43, Cambridge University Press, Cambridge, 1995.

[17] D. Diniz, G. A. Muñoz-Fernández, D. Pellegrino, and J. B. Seoane-Sepúlveda, *The asymptotic growth of the constants in the Bohnenblust–Hille inequality is optimal*, J. Funct. Anal. **263** (2012), 415–428.

[18] ———, *Lower bounds for the constants in the Bohnenblust–Hille inequality: the case of real scalars*, Proc. Amer. Math. Soc., accepted for publication.

[19] P. Erdös, *Some remarks on polynomials*, Bull. Amer. Math. Soc. **53** (1947), 1169–1176.

[20] J. L. Gámez-Merino, G. A. Muñoz-Fernández, V. Sánchez, and J. B. Seoane-Sepúlveda, *Inequalities for polynomials on the unit square via the Krein–Milman Theorem*, J. Convex Analysis **20** (2013), no. 1.

[21] S. Kaijser, *Some results in the metric theory of tensor products*, Studia Math. **63** (1978), no. 2, 157–170.

[22] P. Kirwan, *Complexification of multilinear mappings and polynomials*, Math. Nachr. **231** (2001), 39–68.

[23] M. Krein and D. Milman, *On extreme points of regular convex sets*, Studia Math. **9** (1940), 133–138.

[24] J. E. Littlewood, *On bounded bilinear forms in an infinite number of variables*, Q. J. Math. **1** (1930), 164–174.

[25] A. Montanaro, *Some applications of hypercontractive inequalities in quantum information theory*, J. Math. Physics, accepted for publication.

[26] G. A. Muñoz-Fernández, D. Pellegrino, and J. B. Seoane-Sepúlveda, *Estimates for the asymptotic behavior of the constants in the Bohnenblust–Hille inequality*, Linear Multilinear Algebra **60** (2012), no. 5, 573–582.

[27] G. A. Muñoz-Fernández, S. Gy. Révész, and J. B. Seoane-Sepúlveda, *Geometry of homogeneous polynomials on non symmetric convex bodies*, Math. Scand. **105** (2009), no. 1, 147–160.

[28] G. A. Muñoz, Y. Sarantopoulos, and A. Tonge, *Complexifications of real Banach spaces, polynomials and multilinear maps*, Studia Math. **134** (1999), no. 1, 1–33.

[29] G. A. Muñoz-Fernández and J. B. Seoane-Sepúlveda, *Geometry of Banach spaces of trinomials*, J. Math. Anal. Appl. **340** (2008), no. 2, 1069–1087.

[30] D. Nuñez-Alarcón, D. Pellegrino, and J. B. Seoane-Sepúlveda, *On the Bohnenblust–Hille inequality and a variant of Littlewood's 4/3 inequality*, J. Funct. Anal. **264** (2013), 326–336.

[31] D. Nuñez-Alarcón, D. Pellegrino, J. B. Seoane-Sepúlveda, and D. M. Serrano-Rodríguez, *There exist multilinear Bohnenblust–Hille constants* $(C_n)_{n=1}^{\infty}$ *with* $\lim_{n\to\infty} C_{n+1} - C_n = 0$, J. Funct. Anal. **264** (2013), 429–464.

[32] D. Pellegrino and J. B. Seoane-Sepúlveda, *New upper bounds for the constants in the Bohnenblust–Hille inequality*, J. Math. Anal. Appl. **386** (2012), no. 1, 300–307.

[33] H. Queffélec, *H. Bohr's vision of ordinary Dirichlet series: old and new results*, J. Anal. **3** (1995), 43–60.

[34] H.-J. Rack, *A generalization of an inequality of V. Markov to multivariate polynomials*, J. Approx. Theory **35** (1982), no. 1, 94–97.

[35] J. Siciak, *Wiener's type sufficient conditions for regularity in* $\mathbf{C}^N$, Complex analysis and geometry (Paris, 1997), Progr. Math., vol. 188, Birkhäuser, Basel, 2000, pp. 39–46.

[36] C. Visser, *A generalization of Tchebychef's inequality to polynomials in more than one variable*, Nederl. Akad. Wetensch., Proc. **49** (1946), 455–456 = Indagationes Math. 8, 310–311 (1946).

Departamento de Ciências Exatas,
Universidade Federal da Paraíba,
58.297-000 - Rio Tinto, Brazil.
E-mail address: `jamilson@dce.ufpb.br and jamilsonrc@gmail.com`

Departamento de Análisis Matemático,
Facultad de Ciencias Matemáticas,
Plaza de Ciencias 3,
Universidad Complutense de Madrid,
Madrid, 28040, Spain.
E-mail address: `gustavo_fernandez@mat.ucm.es`

Departamento de Matemática,
Universidade Federal da Paraíba,
58.051-900 - João Pessoa, Brazil.
E-mail address: `pellegrino@pq.cnpq.br and dmpellegrino@gmail.com`

Departamento de Análisis Matemático,
Facultad de Ciencias Matemáticas,
Plaza de Ciencias 3,
Universidad Complutense de Madrid,
Madrid, 28040, Spain.
E-mail address: `jseoane@mat.ucm.es`